

SYNCHRONIZED RF & HIFEM: FAT HISTOLOGY & SCANNING ELECTRON MICROSCOPY STUDY

SIMULTANEOUS APPLICATION OF HIFEM AND SYNCHRONIZED RADIOFREQUENCY FOR FAT DISRUPTION: HISTOLOGICAL AND ELECTRON MICROSCOPY PORCINE MODEL STUDY


Robert A. Weiss MD, FAAD¹, MVDr. Jan Bernardy Ph.D.²,
Prof. MVDr. Frantisek Tichy, CSc³

1. Maryland Laser Skin & Vein Institute, Hunt Valley, MD, USA; 2. Veterinary Research Institute, Brno, CZ
3. Department of Anatomy and Histology, University of Veterinary and Pharmaceutical Sciences Brno, CZ

Presented at the Annual Meeting of the American Society for Dermatologic Surgery, 2020 Virtual Meeting.

HIGHLIGHTS

- Both **histology** and **scanning electron microscopy** showed **damaged adipocytes** post-treatment due to apoptosis and lipolysis.
- **Adipocyte size** was **decreased by 31.1%** at 2 weeks post-treatment.
- The **temperature** in fat tissue was maintained **just below 45°C** for the entire treatment.
- **No necrosis** was seen in the tissue.




Healthy fat cells with well-defined shape at the baseline (left); shrunk adipocytes with noticeable membrane ruptures occurred at 4 days (center); disrupted adipocytes with extrusion of lipid droplets at two weeks (right)

STUDY DESIGN

- 7 Large White pigs (approximately 6 months old).
- All animals received three 30-minute treatments applied to abdomen.
- Biopsy specimens of fat tissue were collected at baseline, 4 days, 2 weeks, 1 month and 2 months post-treatment for each animal.
- Control specimens were collected from the site opposite to the treatment site.
- Evaluation included scanning electron microscopy and histology.

CONCLUSION

- The procedure elevates the **temperature** in subcutaneous fat to levels **necessary** for **apoptosis induction**.
- **Efficacy** of the procedure for **disruption of adipocytes** was documented in **252** analyzed tissue slices.
- Mild inflammatory response was present to promote the **apoptotic death cells removal**.
- The procedure was **safe, no burns, no necrosis** or other adverse events were documented.



Baseline histology (left) showed adipocytes without any damage. At 2 weeks (right), flattened adipocytes with delaminated membranes are seen along with immune cells clearing the damaged tissue.