

Dr. Anil Dalvi and Dr. Matt Mosseler 40 Hillside Drive, Unit 204 Georgetown ON L7G 4W4 Phone: 905-702-2629 Fax: 905-702-5225 info@georgetowngeneralsurgery.com

# COLONOSCOPY – INSTRUCTIONS

# Date of Procedure:

# **Arrival Time:**

### **Procedure Time:**

\*\* Note cancellation or rescheduling within 1 week of the procedure may incur a \$100 cancellation fee at the discretion of the surgeon

## **Location & Details:**

Georgetown Hospital. 1 Princess Anne Drive, Georgetown Ontario. L7G 2B8. Paid Parking is available at the front of the Hospital. Take ticket at parking entrance. Pay machines are located at the entrance of the hospital. Go to the Day Surgery Clinic for registration.

# **Home Medications on day of procedure:**

You should take all your usual morning medications with sips of water (except diabetic pills). If you are
diabetic please discuss your medications with me. Continue your Aspirin or Blood Thinners (e.g.
Coumadin/Warfarin, Plavix) as usual unless instructed otherwise.
Other Instructions:

#### What to bring:

Please bring your Consent Form, Pre-op Surgical Questionnaire, List of home medications, Health card.

#### **After the Procedure:**

After procedure you will be alert once the immediate effects of the sedation wears off over one hr. You will be discharged home with an information sheet indicating the results of your procedure. If any significant results I will speak with you or a designated family member immediately after the procedure.

If you received sedation for your procedure you will need someone to drive you home. You cannot operate a motor vehicle or heavy machinery until the morning after the procedure. If you are taking a taxi home, you MUST have someone accompany you.

You should not drink alcohol or use sleeping pills or other sedatives pills until the following day. <u>Once you depart the hospital there are no other restrictions to your diet and you can restart your home medications unless directed otherwise.</u>

If you notice your abdominal pain or discomfort is worsening after your reach home, or you develop excessive bleeding from your rectum, fevers, or persistent nausea and vomiting go directly to your nearest emergency department.

# **Bowel Preparation**

One box of **Pico-Salax** (2 pouches) **3 Bisacodyl** (Ducolax) **tablets** 

# 3 Days before your procedure

Do NOT eat foods containing seeds, nuts, or corn, raw/uncooked vegetables, potato skins, fruit with skin or seeds until after your procedure. Stop all fibre supplements (flax seeds or Metamucil)

# 1 Day before Colonoscopy

Before 8 am – have a very light, fat free, low fibre breakfast (ex. Toast, bagel, waffle)

After your breakfast you may only drink clear fluids. Aim for a minimum of 3 litres of clear fluids outside of the prep during the day.

Clear Fluids Includes: water, apple juice, consommé/broth, Jell-O, white grape/cranberry juice, black tea (without milk), ginger ale, soda water. Sports drinks like Gatorade are recommended because they contain electrolytes and prevent dehydration. Safest colours are yellow and green.

Do not drink coffee, dairy products or anything colored red, purple or blue.

Diabetics – continue your medication and take 1 meal replacement (such as Glucerna® or Resource®) at each meal today.

## At 4:30 pm take 2 tablets of **Bisacodyl**.

At 5:00 pm drink one packet of **Pico-Salax** in a large glass of water (150 mL). Stir until it has completely dissolved and drink the entire solution. You should drink at least 3 litres of fluid over the next four hours.

Your bowels will usually start to move within 1 hour, producing a series of watery bowel movements. Ensure you are close to a toilet at this time.

## **Day of your Colonoscopy**

At 5:00 am take 1 tablet of **Bisacodyl** and drink the second packet of the **Pico-Salax** solution as above and drink at least 1.5 litres until 2 hours prior to your appointment time. Then nothing to eat or drink. You may have your AM pills with sips as above.

# **Hints for Success**

To lessen the taste of the prep liquid, try drinking it with a straw. Try it cold. To lessen the taste between glasses, try gum or hard candy. For a sore bottom, clean with baby wipes and put on a protective ointment (like Vaseline or Zinc Oxide Cream) after bowel movements. For chills, use a blanket warmed in the dryer.