

# HIFEM TREATMENT OF URINARY INCONTINENCE (UI) IN WOMEN POST GENITOURINARY SURGERY

## HIFEM PROCEDURE FOR TREATMENT OF PERSISTENT URINARY INCONTINENCE POST PELVIC ORGAN PROLAPSE AND ANTI-INCONTINENCE SURGERY


Manish Gopal, M.D., MSCE<sup>1</sup>

1. Center for Urogynecology and Reconstructive Pelvic Surgery, NJ, USA

Presented at 41st Annual Meeting of the American Society for Laser Medicine and Surgery, 2022; San Diego, CA

### HIGHLIGHTS

- Fifty females (60.1±12.3 years), with **pelvic organ prolapse (POP)** and **anti-UI surgery**, received six HIFEM procedures for UI treatment.
- The combined **BFLUTS-SF score decreased significantly** ( $p<0.05$ ) at 3 and 6 months to 11.9±10.2 (-5.5 points) and 12.6±11.7 (-4.8 points).
- The greatest decrease was observed in the **Incontinence domain (-2.0 points,  $p<0.001$ )**, Filling domain (-1.4 points,  $p<0.001$ ) and QoL domain (-1.5 points,  $p<0.01$ ).
- There was a significant improvement in filling, voiding, and incontinence domains regardless of the severity of prolapse.



A decrease in BFLUTS score after the treatment and at all follow-up visits including a linear trendline.