

HIFEM FOR FEMALE SEXUAL DYSFUNCTION: 3-MONTH DATA

THE USE OF HIFEM TECHNOLOGY IN THE TREATMENT OF PELVIC FLOOR MUSCLES AS A CAUSE OF FEMALE SEXUAL DYSFUNCTION: A MULTI-CENTER PILOT STUDY

Timothy Charles Hlavinka MD¹, Pavel Turčan MD², Alexander Bader MD, FAAOCG, FAAC³

1. The Urology Place, InVita Clinics, San Antonio USA

2. Center Mediol, Olomouc, Czech Republic

3. Bader Medical Institute of London, Marylebone, London, United Kingdom


Published in Journal of Women's Health Care: March 9, 2019, DOI: 10.4172/2167-0420.1000455

HIGHLIGHTS


- **30 women** (average age 36±6 years) who experienced challenges related to arousal, achieving orgasm, and painful intercourse were recruited.

Results at 3 months:

- Sexual dysfunction symptoms (FSFI score) **improved** on average by **51%**.
- The most prominent improvement was seen in sexual satisfaction and desire, with a **76%** increase.
- Arousal demonstrated a significant increase, rising by **54%**.
- Patients reported a notable improvement of **60%** in the domain of Orgasm.



SEXUAL SATISFACTION
AND DESIRE



AROUSAL



ORGASM