



ORDER: [REDACTED]
PATIENT: Jennifer Burghart
ID: [REDACTED]
SEX: Female
AGE: [REDACTED]



Microbiology Profile

BACTERIOLOGY CULTURE

Expected/Beneficial flora

4+ *Bacteroides* family
4+ *Bifidobacterium* family
3+ *Escherichia coli*
NG *Lactobacillus* family
4+ *Enterococcus* family
2+ *Clostridium* family

Commensal (Imbalanced) flora

1+ *Citrobacter freundii* complex
2+ *Enterobacter cloacae* complex

Dysbiotic flora

3+ *Klebsiella oxytoca*

NG = No Growth

BACTERIA INFORMATION

Expected / Beneficial bacteria make up a significant portion of the total microflora in a healthy & balanced GI tract. These beneficial bacteria have many health-protecting effects in the GI tract including manufacturing vitamins, fermenting fibers, digesting proteins and carbohydrates, and propagating anti-tumor and anti-inflammatory factors.

Clostridia are prevalent flora in a healthy intestine. *Clostridium* spp. should be considered in the context of balance with other expected/beneficial flora. Absence of clostridia or over abundance relative to other expected/beneficial flora indicates bacterial imbalance. If *C. difficile* associated disease is suspected, a *Clostridium difficile* toxins A/B or GI Pathogens PCR test is recommended.

Commensal (Imbalanced) bacteria are usually neither pathogenic nor beneficial to the host GI tract. Imbalances can occur when there are insufficient levels of beneficial bacteria and increased levels of commensal bacteria. Certain commensal bacteria are reported as dysbiotic at higher levels.

Dysbiotic bacteria consist of known pathogenic bacteria and those that have the potential to cause disease in the GI tract. They can be present due to a number of factors including: consumption of contaminated water or food, exposure to chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels. *Aeromonas*, *Campylobacter*, *Plesiomonas*, *Salmonella*, *Shigella*, *Vibrio*, *Yersinia*, & *Edwardsiella tarda* have been specifically tested for and found absent unless reported.

YEAST CULTURE

Normal flora

No yeast isolated

Dysbiotic flora

MICROSCOPIC YEAST

RESULT

Rare

EXPECTED

None – Rare

Yeast in stool is expected at a level of none-rare. A microscopic finding of yeast in stool of few, moderate, or many may be helpful in identifying potential yeast overgrowth, or non-viable or dietary yeast.

YEAST INFORMATION

Yeast may normally be present in small quantities in the skin, mouth, and intestine. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast are not uniformly dispersed throughout the stool and this may lead to undetectable or low levels of yeast identified by microscopy, despite culture and identified yeast species. Conversely, microscopic examination may reveal a significant amount of yeast present but no viable yeast cultured. Yeast may not always survive transit through the intestines. Nonviable diet-derived yeast may also be detected microscopically. Consideration of clinical intervention for yeast detected microscopically should be made in the context of other findings and presentation of symptoms.

SPECIMEN DATA

Comments:

Date Collected: 08/08/2023

Date Received: 08/10/2023

Date Reported: 08/17/2023

Methodology: Culture and identification by MALDI-TOF and conventional biochemicals





ORDER: [REDACTED]
PATIENT: Jennifer Burghart
ID: [REDACTED]
SEX: Female
AGE: [REDACTED]

CLIENT #: [REDACTED]

Bacterial Susceptibilities; *Klebsiella oxytoca*

NATURAL ANTIBACTERIALS

	LOW SENSITIVITY	HIGH SENSITIVITY	Natural antibacterial agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.
Black Walnut*			
Caprylic Acid*			
Uva Ursi*			
Olive Leaf Extract*			
Oregano*			
Goldenseal*			
Ionic Silver*			
Colloidal Silver*			

PRESCRIPTIVE AGENTS

	RESISTANT	INTERMEDIATE	SUSCEPTIBLE	Susceptible results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used. Intermediate results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used. Resistant results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.
Amoxicillin-Clavulanic Acid			✓	
Ampicillin	✓			
Cefazolin	✓			
Ceftazidime			✓	
Ciprofloxacin			✓	
Sulfamethoxazole / Trimethoprim			✓	

SPECIMEN DATA

Comments:

Date Collected: 08/08/2023

Date Received: 08/10/2023

Date Reported: 08/17/2023

Methodology: Disk Diffusion

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Introduction

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific commentaries are presented. If no significant abnormalities are found, commentaries are not presented.

Microbiology

Beneficial Flora

One or more of the expected or beneficial bacteria are low in this specimen. Normally abundant bacteria include *Lactobacillus* spp, *Bifidobacteria* spp, *Clostridium* spp, *Bacteroides fragilis* group, *Enterococcus* spp, and *Escherichia coli*. The beneficial flora have many health-protecting effects in the gut, and as a consequence, are crucial to the health of the whole organism. Some of the roles of the beneficial flora include digestion of proteins and carbohydrates, manufacture of vitamins and essential fatty acids, increase in the number of immune system cells, break down of bacterial toxins and the conversion of flavonoids into anti-tumor and anti-inflammatory factors. *Lactobacilli*, *bifidobacteria*, *clostridia*, and *enterococci* secrete lactic acid as well as other acids including acetate, propionate, butyrate, and valerate. This secretion causes a subsequent decrease in intestinal pH, which is crucial in preventing an enteric proliferation of microbial pathogens, including bacteria and yeast. Many GI pathogens thrive in alkaline environments. *Lactobacilli* also secrete the antifungal and antimicrobial agents lactocidin, lactobacillin, acidolin, and hydrogen peroxide. The beneficial flora of the GI tract have thus been found useful in the inhibition of microbial pathogens, prevention and treatment of antibiotic associated diarrhea, prevention of traveler's diarrhea, enhancement of immune function, and inhibition of the proliferation of yeast.

In a healthy balanced state of intestinal flora, the beneficial bacteria make up a significant proportion of the total microflora. Healthy levels of each of the beneficial bacteria are indicated by either a 2+, 3+ or 4+ (0 to 4 scale). However, in some individuals there is an imbalance or deficiency of beneficial flora and an overgrowth of non-beneficial (imbalance) or even pathogenic microorganisms (dysbiosis). This can be due to a number of factors including: consumption of contaminated water or food; daily exposure of chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

A number of toxic substances can be produced by the dysbiotic bacteria including amines, ammonia, hydrogen sulfide, phenols, and secondary bile acids which may cause inflammation or damage to the brush border of the intestinal lining. If left unchecked, long-term damage to the intestinal lining may result in leaky gut syndrome, fatigue, chronic headaches, and sensitivities to a variety of foods. In addition, pathogenic bacteria can cause acute symptoms such as abdominal pain, nausea, diarrhea, vomiting and fever in cases of food poisoning.

Antibacterial and antifungal susceptibility testing to a variety of prescriptive and natural agents may be provided for the pathogenic organisms that are cultured from this patient's specimen. This testing is intended to provide the practitioner with useful information to help plan an appropriate treatment regimen. A comprehensive program may be helpful in individuals in whom a dysbiotic condition has caused extensive GI damage.

Note: Not all genera or species can be tested for susceptibilities in the laboratory due to their specific growth requirements. In addition, the Centers for Disease Control and Prevention recommend not testing certain organisms such as those associated with food poisoning. If a practitioner has specific questions, please contact customer service.

Imbalanced Flora

Imbalanced flora are those bacteria that reside in the host gastrointestinal tract and neither injure nor benefit the host. Certain dysbiotic bacteria may appear under the imbalanced category if found at low levels because they are not likely pathogenic at the levels detected. Imbalanced bacteria are commonly more abundant in association with insufficiency dysbiosis, and/or a fecal pH more towards the alkaline end of the reference range (5.8 - 7.0). Treatment with antimicrobial agents is unnecessary unless bacteria appear under the dysbiotic category.

Pathogenic/Dysbiotic Flora

In a healthy balanced state of intestinal flora, the beneficial bacteria make up a significant proportion of the total microflora. However, in many individuals there is an imbalance or deficiency of beneficial flora (insufficiency dysbiosis) and an overgrowth of non-beneficial (imbalance) or even pathogenic microorganisms. This can be due to a number of factors including: consumption of contaminated water or food; daily exposure of chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

A number of toxic substances can be produced by the dysbiotic bacteria including amines, ammonia, hydrogen sulfide, phenols, and secondary bile acids which may cause inflammation or damage to the brush border of the intestinal lining. If left unchecked, long-term damage to the intestinal lining may result in leaky gut syndrome, allergies, autoimmune disease (e.g. rheumatoid arthritis), irritable bowel syndrome, fatigue, chronic headaches, and sensitivities to a variety of foods. In addition, pathogenic bacteria can cause acute symptoms such as abdominal pain, nausea, diarrhea, vomiting, and fever in cases of food poisoning.

Microbiology continued...

Bacterial sensitivities to a variety of prescriptive and natural agents have been provided for the pathogenic bacteria that were cultured from this patient's specimen. This provides the practitioner with useful information to help plan an appropriate treatment regimen. Supplementation with probiotics or consumption of foods (yogurt, kefir, miso, tempeh, tamari sauce) containing strains of lactobacilli, bifidobacteria, and enterococci may help restore healthy flora levels. Soluble fiber and polyphenols derived from chocolate, green tea, blackcurrant, red wine and grape seed extracts have been found to increase the numbers of beneficial bacteria. Hypochlorhydria may also predispose an individual to bacterial overgrowth, particularly in the small intestine. Nutritional anti-inflammatories can aid in reversing irritation to the GI lining. These include quercetin, vitamin C, curcumin, gamma-linoleic acid, omega-3 fatty acids (EPA, DHA), and aloe vera. Other nutrients such as zinc, beta-carotene, pantothenic acid, and L-glutamine provide support for regeneration of the GI mucosa. A comprehensive program may be helpful in individuals in whom a dysbiotic condition has caused extensive GI damage.

***Klebsiella* spp**

Klebsiella spp. are gram-negative bacilli belonging to the *Enterobacteriaceae* family and closely related to the genera *Enterobacter* and *Serratia*. *Klebsiella* spp. are considered dysbiotic in the amount of 3 - 4 +. *Klebsiella* spp. are widely distributed in nature and in the gastrointestinal tract of humans. In humans, they may colonize the skin, oral cavity, pharynx, or gastrointestinal tract. Regarded as normal flora in many parts of the colon, intestinal tract and biliary tract, the gut is the main reservoir of opportunistic strains. This bacteria has the potential to cause intestinal, lung, urinary tract, and wound infections, but overgrowth of *Klebsiella* spp. is commonly asymptomatic. *K. pneumoniae*, in particular, may cause diarrhea and some strains are enterotoxigenic. Infection has been linked to ankylosing spondylitis as well as myasthenia gravis (antigenic cross-reactivity), and these patients usually carry larger numbers of the organism in their intestines than healthy individuals. *Klebsiella oxytoca* causes antibiotic associated hemorrhagic colitis. These strains have been shown to produce a cytotoxin that is capable of inducing cell death in various epithelial-cell cultures.

Klebsiella is a significant nosocomial infectious agent, partially due to the ability of organisms to spread rapidly. *Klebsiella* accounts for approximately 3-7% of all hospital-acquired infections, placing it among the top eight pathogens in hospitals. Extraintestinal infection typically involves the respiratory or urinary tracts, but may infect other areas such as the biliary tract and surgical wound sites. *K. pneumoniae* and *K. oxytoca* are the two members of this genus responsible for most extraintestinal human infections.

Treatment of these organisms has become a major problem because of resistance to multiple antibiotics and potential transfer of plasmids to other organisms. Proper hand washing is crucial to prevent transmission from patient to patient via medical personnel. Contact isolation should be used for patients colonized or infected with highly antibiotic-resistant *Klebsiella* strains. *Klebsiella ozaenae* and *Klebsiella rhinoscleromatis* are infrequent isolates that are subspecies of *K. pneumoniae*; however, each is associated with a unique spectrum of disease. *K. ozaenae* is associated with atrophic rhinitis, a condition called ozena, and purulent infections of the nasal mucous membranes. *K. rhinoscleromatis* causes the granulomatous disease rhinoscleroma, an infection of the respiratory mucosa, oropharynx, nose, and paranasal sinuses.

Antibiotics may be indicated if symptoms are prolonged and in systemic infections. Refer to the antimicrobial susceptibilities for treatment.